

Superfluidity of “dirty” indirect magnetoexcitons in coupled quantum wells in high magnetic field

Oleg L. Berman¹, Yurii E. Lozovik², David W. Snoke³, and Rob D. Coalson¹

¹ *Department of Chemistry, University of Pittsburgh,
Pittsburgh, PA 15260, USA*

² *Institute of Spectroscopy, Russian Academy of Sciences,
142190 Troitsk, Moscow Region, Russia*

³ *Department of Physics and Astronomy, University of Pittsburgh,
Pittsburgh, PA 15260 USA*

Abstract

Superfluidity in the quasi-two-dimensional (2D) system of spatially indirect magnetoexcitons in coupled quantum wells (CQW) and unbalanced two-layer electron system in high magnetic field H is considered in the presence of a random field. The problem of the rare gas of magnetoexcitons with dipole-dipole repulsion in a random field has been reduced to the problem of the rare gas of dipole excitons without magnetic field with the effective magnetic mass of a magnetoexciton, which is a function of the magnetic field and parameters of the CQW, in an H -dependent effective random field. The density of the superfluid component n_s and the temperature T_c of the Kosterlitz-Thouless transition to a superfluid state are obtained as functions of magnetic field H , interlayer separation D and the random field parameters α_i and g_i . For 2D magnetoexcitonic systems, the rise of the magnetic field H and the interwell distance D is found to increase the effective renormalized random field parameter Q and suppress the superfluid density n_s and the temperature of the Kosterlitz-Thouless transition T_c . The suppressing influence of D on n_s and T_c in strong magnetic field is opposite to the case without magnetic field, when n_s and T_c increase with the rise of D at fixed total exciton density n . It is shown that in the presence of the disorder at sufficiently large magnetic field H or parameters of the disorder there is no superfluidity at any exciton density.

Key words: coupled quantum wells (CQW), nanostructures, superfluidity, magnetoexciton, Bose-Einstein condensation of magnetoexcitons.

PACS numbers: 73.20.Dx, 71.35.Ji, 71.35.Lk

I. INTRODUCTION

Systems of excitons with spatially separated electrons (e) and holes (h) (indirect excitons) in coupled quantum wells (CQW) in magnetic fields (H) have been the subject of several experimental investigations.^{1–7} These systems without magnetic field were also under the consideration recently.^{8,9} They are of interest, in particular, in connection with the possibility of superfluidity of indirect excitons or $e-h$ pairs, which would manifest itself in the CQW as persistent electrical currents in each well^{10–17}. In high magnetic fields ($H > 7T$) two-dimensional (2D) excitons survive in a substantially wider temperature region, as the exciton binding energies increase with magnetic field.^{18–27} In addition, the 2D $e-h$ system in high H fields is interesting due to the existence, under some conditions of supersymmetry in the system (for the single quantum well) leading to unique exact solutions of the many-body problem (the last corresponding to ideal Bose-condensation of magnetoexcitons at *any* density)^{18,20}.

The superfluid state appears in the system under consideration below the temperature of Kosterlitz-Thouless transition.²⁸ The latter was studied recently for systems with spatially separated electrons (e) and holes (h) in the absence of magnetic field.^{16,17}

Attempts at experimental investigation of magnetoexciton superfluidity in coupled quantum wells¹ make it essential to study the magnetic field dependence of the temperature of phase transition to the superfluid state in systems of indirect magnetoexcitons and to analyze the density of the superfluid component. It was shown that increasing the magnetic field at a fixed magnetoexciton density leads to a lowering of the Kosterlitz-Thouless transition temperature T_c on account of the increase of the exciton magnetic mass as a function of a magnetic field H ²⁵. But it turns out that the highest possible Kosterlitz-Thouless transition temperature increases with H (at small D) due to the rise in the maximal density of magnetoexcitons *vs.* H ²⁵.

The theory of the magnetoexcitonic systems did not take into account the role of disorder, which is created by impurities and boundary irregularities of the quantum wells. In real experiments, however, disorder plays a very important role. Although the inhomogeneous broadening linewidth of typical GaAs-based samples has been improved from around 20 meV to less than 1 meV,⁸ the disorder energy is still not much smaller than the exciton-exciton repulsion energy. At a typical exciton density of 10^{10} cm^{-2} , the interaction energy of the

excitons is approximately 1 meV.⁷ On the other hand, the typical disorder energy of 1 meV is low compared to the typical exciton binding energy of 5 meV.

A typical example of a two-dimensional system of weakly interacting bosons is a system of indirect excitons in coupled quantum wells (GaAs/AlGaAs)^{1,3,4,5,6,8,9}. Fluctuations of the thickness of a quantum well, which arise during the fabrication process, impurities in the system, and disorder in the alloy of the barriers can all lead to the appearance of a random field. Of these, spectral analysis of the exciton luminescence shows that alloy disorder, with a characteristic length scale short compared to the excitonic Bohr radius of around 100 Å, plays the most important role.

The collective properties of 2D indirect excitons in weak disorder without magnetic field were analyzed in Ref.[29]. In Ref.[30] we studied the case of a random field which is not necessarily small compared to the dipole-dipole repulsion between excitons for CQW without magnetic field. The coherent potential approximation (CPA) allows us to derive the 2D indirect exciton Green's function for a wide range of the random field, resulting in the second order Born approximation in weak scattering limit (the second order Born approximation Green's function for 3D excitons was obtained by Gevorkyan and Lozovik³¹). It was predicted that in the low-temperature limit, the density n_s of the superfluid component in CQW systems and the temperature of the superfluid transition (the Kosterlitz-Thouless temperature T_c ²⁸) are the decreasing functions of the random field³⁰.

In Ref.[32] the spectrum of the single indirect 2D magnetoexciton (non-interacting with other magnetoexcitons) in the strong perpendicular magnetic field in the presence of disorder in the second-order Born approximation was obtained.

In the present paper we derive the collective spectrum of the 2D low-dense indirect magnetoexciton gas in the presence of the disorder including the dipole-dipole repulsion between magnetoexcitons in the ladder approximation²⁵. We consider disorder which is not weak compared to the dipole repulsion in the second-order Born approximation³². The density of the superfluid component n_s and the temperature T_c of the Kosterlitz-Thouless transition to a superfluid state are obtained as a function of magnetic field H , interlayer separation D and the random field parameters α_i and g_i . We show that the density of the superfluid component and the Kosterlitz-Thouless temperature of the superfluid phase transition decrease as the random field increases. These results are derived by the mapping the problem of the rare indirect magnetoexciton gas with the disorder in the strong magnetic

field onto the problem of the rare dipole gas without magnetic field consisting of indirect excitons with the effective magnetic mass of a magnetoexciton, which is a function of the magnetic field and the parameters of the coupled quantum wells (CQW), in an H -dependent effective random field. The application of these results, obtained for CQW with spatially separated electrons and holes, for unbalanced two-layer electron system is also discussed.

The paper is organized in the following way. In Sec. II we derive the effective Hamiltonian of rare “dirty” spatially indirect magnetoexcitons in the effective magnetic mass approximation in high magnetic field limit. In Sec. III we obtain the Green’s function of a single magnetoexciton in the random field. In Sec. IV the collective spectrum, the superfluid density and the temperature of the Kosterlitz-Thouless phase transition of magnetoexcitons are derived in the presence of disorder. In Sec. V we discuss our results and consider the extension of the framework used for the indirect magnetoexcitons in CQW on the system of indirect magnetoexcitons in unbalanced two-layer electron system.

II. THE EFFECTIVE HAMILTONIAN OF RARE “DIRTY” SPATIALLY INDIRECT MAGNETOEXCITONS IN THE EFFECTIVE MAGNETIC MASS APPROXIMATION

The total Hamiltonian $\hat{H}$ of 2D spatially separated e and h in the perpendicular magnetic field in the presence of the external field in the second quantization representation has the form:

$$\begin{aligned} \hat{H} = & \int d\mathbf{R} \int d\mathbf{r} \left[\hat{\psi}^\dagger(\mathbf{R}, \mathbf{r}) \left(\frac{1}{2m_e} (-i\nabla_e + e\mathbf{A}_e)^2 + \frac{1}{2m_h} (-i\nabla_h - e\mathbf{A}_h)^2 \right. \right. \\ & \left. \left. - \frac{e^2}{\epsilon\sqrt{(\mathbf{r}_e - \mathbf{r}_h)^2 + D^2}} + V_e(\mathbf{r}_e) + V_h(\mathbf{r}_h) \right) \hat{\psi}(\mathbf{R}, \mathbf{r}) \right] \\ & + \frac{1}{2} \int d\mathbf{R}_1 \int d\mathbf{r}_1 \int d\mathbf{R}_2 \int d\mathbf{r}_2 \hat{\psi}^\dagger(\mathbf{R}_1, \mathbf{r}_1) \hat{\psi}^\dagger(\mathbf{R}_2, \mathbf{r}_2) \left(U^{ee}(\mathbf{r}_{e1} - \mathbf{r}_{e2}) + U^{hh}(\mathbf{r}_{h1} - \mathbf{r}_{h2}) \right. \\ & \left. + U^{eh}(\mathbf{r}_{e1} - \mathbf{r}_{h2}) + U^{he}(\mathbf{r}_{h1} - \mathbf{r}_{e2}) \right) \hat{\psi}(\mathbf{R}_2, \mathbf{r}_2) \hat{\psi}(\mathbf{R}_1, \mathbf{r}_1), \end{aligned} \quad (1)$$

Here $\hat{\psi}^\dagger(\mathbf{R}, \mathbf{r})$ and $\hat{\psi}(\mathbf{R}, \mathbf{r})$ are the operators of creation and annihilation of magnetoexcitons; $\mathbf{r}_e$ and $\mathbf{r}_h$ are electron and hole locations along quantum wells, correspondingly; $\mathbf{A}_e$, $\mathbf{A}_h$ are the vector potentials in electron and hole location, correspondingly; $V_e(\mathbf{r}_e)$ and $V_h(\mathbf{r}_h)$ represent the external fields acting on electron and hole, correspondingly (we use units $c = \hbar = 1$); D is the distance between electron and hole quantum wells; e is the charge of an

electron; ϵ is the dielectric constant. We use below the coordinates of the magnetoexciton center of mass $\mathbf{R} = (m_e \mathbf{r}_e + m_h \mathbf{r}_h)/(m_e + m_h)$ and the internal exciton coordinates $\mathbf{r} = \mathbf{r}_e - \mathbf{r}_h$. The cylindrical gauge for vector-potential is used: $\mathbf{A}_{e,h} = \frac{1}{2} \mathbf{H} \times \mathbf{r}_{e,h}$. U^{ee} , U^{hh} , U^{eh} and U^{he} are the two-particle potentials of the electron-electron, hole-hole, electron-hole and hole-electron interaction, correspondingly, between electrons or holes from different pairs:

$$\begin{aligned} U^{ee}(\mathbf{r}_{e1} - \mathbf{r}_{e2}) &= \frac{e^2}{\epsilon |\mathbf{r}_{e1} - \mathbf{r}_{e2}|}; & U^{hh}(\mathbf{r}_{h1} - \mathbf{r}_{h2}) &= \frac{e^2}{\epsilon |\mathbf{r}_{h1} - \mathbf{r}_{h2}|}; \\ U^{eh}(\mathbf{r}_{e1} - \mathbf{r}_{h2}) &= -\frac{e^2}{\epsilon \sqrt{|\mathbf{r}_{e1} - \mathbf{r}_{h2}|^2 + D^2}}; & U^{he}(\mathbf{r}_{h1} - \mathbf{r}_{e2}) &= -\frac{e^2}{\epsilon \sqrt{|\mathbf{r}_{h1} - \mathbf{r}_{e2}|^2 + D^2}}. \end{aligned} \quad (2)$$

The conserved quantity for isolated exciton in magnetic field without any external field ($V_e(\mathbf{r}_e) = V_h(\mathbf{r}_h) = 0$) (the exciton *magnetic* momentum) is (see Ref.[33]):

$$\hat{\mathbf{P}} = -i\nabla_e - i\nabla_h + e(\mathbf{A}_e - \mathbf{A}_h) - e\mathbf{H} \times (\mathbf{r}_e - \mathbf{r}_h), \quad (3)$$

The conservation of this quantity is related to the invariance of the system upon a simultaneous translation of e and h and gauge transformation.

The eigenfunctions of Hamiltonian of a single isolated magnetoexciton without any random field ($V_e(\mathbf{r}_e) = V_h(\mathbf{r}_h) = 0$), which are also the eigenfunctions of the magnetic momentum $\mathbf{P}$, have the following form (see Refs. [18,33]):

$$\Psi_{k\mathbf{P}}(\mathbf{R}, \mathbf{r}) = \exp \left\{ i\mathbf{R} \left(\mathbf{P} + \frac{e}{2} \mathbf{H} \times \mathbf{R} \right) + i\gamma \frac{\mathbf{P}\mathbf{r}}{2} \right\} \Phi_k(\mathbf{P}, \mathbf{r}), \quad (4)$$

where $\Phi_k(\mathbf{P}, \mathbf{r})$ is the function of internal coordinates $\mathbf{r}$; $\mathbf{P}$ is the eigenvalue of magnetic momentum; k are quantum numbers of j exciton internal motion. In high magnetic fields $k = (n_L, m)$, where $n_L = \min(n_1, n_2)$, $m = |n_1 - n_2|$, $n_{1(2)}$ are Landau quantum numbers for e and h ^{18,24}; $\gamma = (m_h - m_e)/(m_h + m_e)$.

In this section we reduce the problem of the rare gas of magnetoexcitons (in high magnetic field) with dipole-dipole repulsion in a random field to the problem of the rare gas of dipole excitons without magnetic field with the effective magnetic mass of a magnetoexciton, which is a function of the magnetic field and parameters of the quantum wells, in an H -dependent effective random field. Our goal is to map the Hamiltonian of the rare gas of “dirty” dipole-dipole repulsing magnetoexcitons placed on the lowest Landau level (high magnetic field regime) onto the the Hamiltonian of the rare gas of “dirty” dipole-dipole repulsing excitons without the magnetic field, but characterized by the effective magnetic mass dependent on H and D and the effective random field, renormalized by magnetic field H .

We further expand the magnetoexciton field operators in a single magnetoexciton basis set $\Psi_{k\mathbf{P}}(\mathbf{R}, \mathbf{r})$

$$\hat{\psi}^\dagger(\mathbf{R}, \mathbf{r}) = \sum_{k\mathbf{P}} \Psi_{k\mathbf{P}}^*(\mathbf{R}, \mathbf{r}) \hat{a}_{k\mathbf{P}}^\dagger; \quad \hat{\psi}(\mathbf{R}, \mathbf{r}) = \sum_{k\mathbf{P}} \Psi_{k\mathbf{P}}(\mathbf{R}, \mathbf{r}) \hat{a}_{k\mathbf{P}}, \quad (5)$$

where $\hat{a}_{k\mathbf{P}}^\dagger$ and $\hat{a}_{k\mathbf{P}}$ are the corresponding creation and annihilation operators of a magnetoexciton in $(k, \mathbf{P})$ space.

We consider the case of strong magnetic field, when we neglect in Eq. (4) the transitions between different Landau levels of the magnetoexciton caused by scattering by the slowly changing in space potential $V_e(\mathbf{r}_e) + V_h(\mathbf{r}_h)$. We also neglect nondiagonal matrix elements of the Coulomb interaction between electron and hole in the same pair. The application region of these two assumptions is defined by the inequalities

$$\omega_c \gg E_b, \quad \omega_c \gg \sqrt{\langle V_{e(h)}^2 \rangle_{av}}, \quad (6)$$

where $\omega_c = \sqrt{eH/m_{e-h}}$, $m_{e-h} = m_e m_h / (m_e + m_h)$ is the exciton reduced mass in the quantum well plane; E_b is the magnetoexciton binding energy in an ideal “pure” system as a function of magnetic field H and the distance between electron and hole quantum wells D : $E_b \sim e^2/\epsilon r_H \sqrt{\pi/2}$ at $D \ll r_H$ and $E_b \sim e^2/\epsilon D$ at $D \gg r_H$ ($r_H = (eH)^{-1/2}$ is the magnetic length).^{18,24} Here $\langle \dots \rangle_{av}$ denotes averaging over the fluctuations of random field.

After exciton-exciton scattering their total magnetic momentum $\mathbf{P} = \sum_i \mathbf{P}_i$ is conserved, but the magnetic momentum $\mathbf{P}_i$ of each exciton can be changed. Since in a strong magnetic field ρ , the mean distance between e and h along the quantum wells is proportional to the magnetic momentum ($\rho = \frac{r_H^2}{H} \mathbf{H} \times \mathbf{P}$)^{18,33}, the scattering is accompanied by the exciton polarization. We consider sufficiently low temperatures when magnetoexciton states with only small magnetic momenta $P \ll \frac{1}{r_H}$ are filled. The change of these magnetic momenta due to exciton-exciton scattering is also negligible due to the conservation of the total magnetic momentum. But these small magnetic momenta correspond to small separation between electrons and holes along quantum wells $\rho \ll r_H$. So magnetoexciton polarization due to scattering is negligible and the magnetoexciton dipole moment keeps to be almost normal to quantum wells $d = eD$, i.e. the interexciton interaction law is not changed due to the scattering. For the rare system in the strong magnetic field $n \ll r_H^{-2}$ (n is the 2D density of excitons; the characteristic radius of magnetoexciton along the wells in the strong magnetic field at small P approximately equals to the magnetic length r_H ²⁴) the exciton-exciton

interaction is the dipole-dipole repulsion, because the average distance between excitons r_s is large compare to the exciton radius ($r_s = (\pi n)^{-1/2} \gg r_H$). So for the rare system in the strong magnetic field at $n \ll r_H^{-2}$ we have:

$$\hat{U}(|\mathbf{R}_1 - \mathbf{R}_2|) \equiv \hat{U}^{ee} + \hat{U}^{hh} + \hat{U}^{eh} + \hat{U}^{he} \simeq \frac{e^2 D^2}{\epsilon |\mathbf{R}_1 - \mathbf{R}_2|^3}. \quad (7)$$

Notice that the magnetoexcitons repel like parallel dipoles, and their pair interaction potential depends only on the coordinates of the center mass of the excitons and does not depend on the coordinates of the relative motion of the electron and hole.

Now we substitute the expansions Eq. (5) for the field creation and annihilation operators in the total Hamiltonian Eq. (1) and obtain the effective Hamiltonian in terms of creation and annihilation operators in $\mathbf{P}$ space. In strong magnetic fields $\omega_c = eH/\mu^* \gg e^2/r_H$ the characteristic value of $e - h$ separation in the magnetoexciton $|\langle \mathbf{r} \rangle|$ has the order of the magnetic length $r_H = 1/\sqrt{eH}$. The functions $\Phi_k(\mathbf{P}, \mathbf{r})$ (see Eq.(4)) are dependent on the difference $(\mathbf{r} - \rho)$, where $\rho = \frac{r_H^2}{H} \mathbf{H} \times \mathbf{P}$ ^{18,33}. At small magnetic momenta $P \ll 1/r_H$ we have $\rho \ll r_H$, and, therefore, in functions $\Phi_k(\mathbf{r} - \rho)$ we can ignore the variable ρ in comparison with $\mathbf{r}$. In strong magnetic fields quantum numbers k correspond to the quantum numbers (m, n_L) (see above). For the lowest Landau level we denote the spectrum of the single exciton $\varepsilon_0(P) \equiv \varepsilon_{00}(\mathbf{P})$. In high magnetic field, when the typical interexciton interaction $D^2 n^{-3/2} \ll \omega_c$, one can ignore transitions between Landau levels and consider only the states on the lowest Landau level $m = n_L = 0$. Using the orthonormality of functions $\Phi_{mn}(\mathbf{0}, \mathbf{r})$ we obtain the effective Hamiltonian $\hat{H}_{eff}$ in strong magnetic fields. Since a typical value of r is r_H , and $P \ll 1/r_H$ in this approximation the effective Hamiltonian $\hat{H}_{eff}$ in the magnetic momentum representation P on the lowest Landau level $m = n_L = 0$ has the same form (compare with Ref.[17]) as for two-dimensional boson system without a magnetic field, but with the magnetoexciton magnetic mass m_H (which depends on H and D) instead of the exciton mass ($M = m_e + m_h$), magnetic momenta instead of ordinary momenta and renormalized random field:

$$\begin{aligned} \hat{H}_{eff} = & \sum_{\mathbf{P}} \varepsilon_0(P) \hat{a}_{\mathbf{P}}^\dagger \hat{a}_{\mathbf{P}} + \sum_{\mathbf{P}, \mathbf{P}'} \langle \mathbf{P}' | \hat{V} | \mathbf{P} \rangle \hat{a}_{\mathbf{P}'}^\dagger \hat{a}_{\mathbf{P}} \\ & + \frac{1}{2} \sum_{\mathbf{P}_1, \mathbf{P}_2, \mathbf{P}_3, \mathbf{P}_4} \langle \mathbf{P}_1, \mathbf{P}_2 | \hat{U} | \mathbf{P}_3, \mathbf{P}_4 \rangle \hat{a}_{\mathbf{P}_1}^\dagger \hat{a}_{\mathbf{P}_2}^\dagger \hat{a}_{\mathbf{P}_3} \hat{a}_{\mathbf{P}_4}, \end{aligned} \quad (8)$$

where $\hat{V} = \hat{V}_e + \hat{V}_h$. The dispersion relation $\varepsilon_0(P)$ of an isolated magnetoexciton on the lowest

Landau level is the quadratic function at small magnetic momenta under consideration:

$$\varepsilon_0(\mathbf{P}) \approx \frac{P^2}{2m_H}, \quad (9)$$

where m_H is the effective *magnetic* mass of a magnetoexciton on the lowest Landau level, dependent on H and the distance D between e – and h – layers (see Ref.[24]). In strong magnetic fields at $D \gg r_H$ the exciton magnetic mass is $m_H \approx D^3\epsilon/(e^2r_H^4)^{24}$. The quadratic dispersion relation Eq.(9) is true for small P at arbitrary magnetic fields H and follows from the fact that $P = 0$ is an extremal point of the dispersion law $\varepsilon_k(\mathbf{P})$. The last statement may be proved by taking into account the regularity of the effective Hamiltonian $H_{\mathbf{P}}$ as a function of the parameter $\mathbf{P}$ at $\mathbf{P} = 0$ and also the invariance of $H_{\mathbf{P}}$ upon simultaneous rotation of $\mathbf{r}$ and $\mathbf{P}$ in the CQW plane. For magnetoexciton ground state $m_H > 0$. For high magnetic fields $r_H \ll a_0^*$ and at $D \lesssim r_H$ the quadratic dispersion relation is valid at $P \ll 1/r_H$, but for $D \gg r_H$ it holds over a wider region — at least at $P \ll D/r_H^2$ ²⁴ ($a_0^* = \epsilon/(2m_{e-h}e^2)$ is the radius of a 2D exciton at $H = 0$).

The matrix element of the inter-magnetoexciton interaction $\langle \mathbf{P}_1, \mathbf{P}_2 | \hat{U} | \mathbf{P}_3, \mathbf{P}_4 \rangle$ is defined as

$$\begin{aligned} \langle \mathbf{P}_1, \mathbf{P}_2 | \hat{U} | \mathbf{P}_3, \mathbf{P}_4 \rangle = & \frac{1}{S^2} \int d^2R_1 \int d^2R_2 \int d^2r_1 \int d^2r_2 U(\mathbf{R}_1 - \mathbf{R}_2) \\ & \Psi_{k_1\mathbf{P}_1}^*(\mathbf{R}_1, \mathbf{r}_1) \Psi_{k_2\mathbf{P}_2}^*(\mathbf{R}_2, \mathbf{r}_2) \Psi_{k_3\mathbf{P}_3}(\mathbf{R}_1, \mathbf{r}_1) \Psi_{k_4\mathbf{P}_4}(\mathbf{R}_2, \mathbf{r}_2). \end{aligned} \quad (10)$$

The matrix potential of $\hat{U}$ (Eq. (7)) connecting the states $\langle k_1 = k_2 = 0, \mathbf{P}_1, \mathbf{P}_2 |$ and $\langle k_3 = k_4 = 0, \mathbf{P}_3, \mathbf{P}_4 |$ has the form

$$\langle \mathbf{P}_1, \mathbf{P}_2 | \hat{U} | \mathbf{P}_3, \mathbf{P}_4 \rangle = \frac{1}{S^2} U(\mathbf{P}_3 - \mathbf{P}_1) \delta(\mathbf{P}_1 + \mathbf{P}_2 - \mathbf{P}_3 - \mathbf{P}_4), \quad (11)$$

where S is the area of a quantum well, and

$$U(\mathbf{P}_3 - \mathbf{P}_1) = \int \int U(|\mathbf{R}_1 - \mathbf{R}_2|) \exp(i(\mathbf{P}_3 - \mathbf{P}_1)(\mathbf{R}_1 - \mathbf{R}_2)) d^2|\mathbf{R}_1 - \mathbf{R}_2|. \quad (12)$$

In the strong magnetic field limit, using for $\Phi_k(\mathbf{P}, \mathbf{r} - \rho)$ ($\rho = \frac{r_H^2}{H} \mathbf{H} \times \mathbf{P}$) in Eq. (4) the internal wavefunction of the magnetoexciton at the lowest Landau level, at small magnetic momenta $P \ll 1/r_H$ we can ignore the variable ρ relatively to $\mathbf{r}$ (see above). So in $\Psi_{k=0,\mathbf{P}}(\mathbf{R}, \mathbf{r})$ (Eq. (4)) we put $\mathbf{P} = 0$ and $\rho = 0$ for $\Phi_{k=0}(\mathbf{P}, \mathbf{r} - \rho)$, while we keep $\mathbf{P}\mathbf{R} \neq 0$ in the exponent. This procedure is valid in a strong magnetic field at small magnetic momenta, because the characteristic ρ is much smaller than the characteristic $R \sim r_s = (\pi n)^{-1/2}$

$(r_H \ll (\pi n)^{-1/2})$. Keeping $\mathbf{P}\mathbf{R} \neq 0$, we can use the magnetic momentum conservation law below. So in the strong magnetic field limit, using for $\Phi_k(\mathbf{0}, \mathbf{r})$ the internal wavefunction of the magnetoexciton at the lowest Landau level²⁴

$$\Phi_{k=0}(\mathbf{0}, \mathbf{r}) = \frac{1}{\sqrt{2\pi}r_H} \exp \left[-\frac{r^2}{4r_H^2} \right], \quad (13)$$

we obtain the matrix element of the external potential $V_{e,h}(\mathbf{r})$ connecting the states $\langle k=0, \mathbf{P} |$ and $\langle k=0, \mathbf{P}' |$, which is defined as

$$\langle \mathbf{P}' | \hat{V}_{e,h}(\mathbf{r}) | \mathbf{P} \rangle = \frac{1}{S} \int d^2R \int d^2r V_{e,h}(\mathbf{r}_{e,h}) \Psi_{k'\mathbf{P}'}^*(\mathbf{R}, \mathbf{r}) \Psi_{k\mathbf{P}}(\mathbf{R}, \mathbf{r}), \quad (14)$$

and has the form

$$\begin{aligned} \langle \mathbf{P}' | \hat{V}_{e,h}(\mathbf{r}) | \mathbf{P} \rangle &= \frac{1}{S} \exp \left(-(\mathbf{P}' - \mathbf{P})^2 \frac{r_H^2}{4} \right) \\ V_{e,h}(\mathbf{P}' - \mathbf{P}) \exp \left(\pm \frac{ir_H^2}{2H} \mathbf{H}\mathbf{P} \times \mathbf{P}' \right) &= \\ = \frac{1}{S} \tilde{V}_{e,h}(\mathbf{P}' - \mathbf{P}) \exp \left(\pm \frac{ir_H^2}{2H} \mathbf{H}\mathbf{P} \times \mathbf{P}' \right), \end{aligned} \quad (15)$$

where

$$V_{e,h}(\mathbf{P}' - \mathbf{P}) = \int \int V_{e,h}(\mathbf{r}) \exp [i(\mathbf{P}' - \mathbf{P})\mathbf{r}] d^2r. \quad (16)$$

Then, using the expressions for the matrix elements Eqs. (10) and (14), the expansions for the the field operators Eq. (5), applying the orthonormality of functions $\Phi_k(\mathbf{0}, \mathbf{r})$, and employing only the lowest Landau level $k=0$ in the strong magnetic field limit at the end we obtain from the Eq. (8) the effective Hamiltonian of dipole indirect magnetoexctions in high magnetic field in the presence of the disorder in coordinate space:

$$\begin{aligned} \hat{H}_{eff} &= \int d\mathbf{R} \hat{\psi}^\dagger(\mathbf{R}) \left(-\frac{\nabla^2}{2m_H} + V_{eff}(\mathbf{R}) \right) \hat{\psi}(\mathbf{R}) \\ &+ \frac{1}{2} \int d\mathbf{R}_1 \int d\mathbf{R}_2 \hat{\psi}^\dagger(\mathbf{R}_1) \hat{\psi}^\dagger(\mathbf{R}_2) U(\mathbf{R}_1 - \mathbf{R}_2) \hat{\psi}(\mathbf{R}_2) \hat{\psi}(\mathbf{R}_1), \end{aligned} \quad (17)$$

where $\hat{\psi}^\dagger(\mathbf{R})$ and $\hat{\psi}(\mathbf{R})$ are the Bose creation and annihilation field operators (the possibility of the assumption about magnetoexcitons being bosons we discuss below), and the coupling to the effective random field $V_{eff}(\mathbf{R})$ has the form

$$V_{eff}(\mathbf{R}) = \frac{1}{\pi r_H^2} \int \exp \left(-\frac{(\mathbf{R} - \mathbf{r})^2}{r_H^2} \right) [V_e(\mathbf{r}) + V_h(\mathbf{r})] d\mathbf{r}. \quad (18)$$

The Eq.(18) is valid, if the correlation length L of random potential ($V(\mathbf{r}_e, \mathbf{r}_h)$) is much greater than the magnetoexciton mean size $r_{exc} = r_H$: ($L \gg r_H$), i.e., in the case of a smooth surface potential, which is realized, as was shown using a scanning tunnelling microscope, on interfaces in $AlGaAs - GaAs$ structures discussed in this paper. A more rigorous condition of the smoothness of random potential can be expressed as³⁴

$$r_{exc} \sqrt{\langle \nabla V^2 \rangle_{av}} \ll E_b, \quad (19)$$

and it holds for the strong magnetic field, when $r_{exc} = r_H = (eH)^{-1/2}$, and $E_b \sim e^2/\epsilon D$ at $D \gg r_H$ ^{24,32}.

As a matter of fact, the effective magnetoexciton Hamiltonian $\hat{H}_{eff}$ (Eq. (17)) treats the magnetoexciton as an electrically neutral composite particle. Since the particle is neutral, it does not directly interact with the magnetic field. The interaction with the magnetic field manifests itself through the renormalization of the exciton effective mass and modification of the correlation function of the random field. Thus, we can do mapping from the original problem of the rare weakly-interacting magnetoexciton system in strong magnetic field in the presence of the disorder, described by the total Hamiltonian Eq. (1), on the rare system of excitons without magnetic field with the effective magnetic mass m_H and in the presence of the effective random field V_{eff} (Eq. (18)), renormalized by the magnetic field H . The dipole-dipole interaction term in the effective Hamiltonian Eq. (17) in the strong magnetic field limit is exactly the same as one for the excitons without magnetic field, because the dipole-dipole interaction does not depend on the relative coordinates of electron and hole in the exciton as long as we assume the excitons are parallel dipoles (as mentioned above). This mapping allows us to use the results for the collective spectrum, superfluid density and Kosterlitz-Thouless temperature, obtained for the system in random field without magnetic field³⁰, for the case of strong magnetic field.

III. THE GREEN'S FUNCTION OF A SINGLE MAGNETOEXCITON IN THE RANDOM FIELD

The interaction between an spatially indirect exciton in coupled quantum wells and a random field, induced by fluctuations in widths of electron and hole quantum wells, has the

form³²:

$$V(\mathbf{r}_e, \mathbf{r}_h) = \alpha_e[\xi_1(\mathbf{r}_e) - \xi_2(\mathbf{r}_e)] + \alpha_h[\xi_3(\mathbf{r}_h) - \xi_4(\mathbf{r}_h)], \quad (20)$$

where $\alpha_{e,h} = \partial E_{e,h}^{(0)} / \partial d_{e,h}$, $d_{e,h}$ is the average widths of the electron and hole quantum wells, $E_{e,h}^{(0)}$ are the lowest levels of the electron and hole in the conduction and valence bands, and ξ_1 and ξ_2 (ξ_3 and ξ_4) are fluctuations in the widths of the electron (hole) wells on the upper and lower interfaces, respectively. We assume that fluctuations on different interfaces are statistically independent, whereas fluctuations of a specific interface are characterized by Gaussian correlation function

$$\langle \xi_i(\mathbf{r}_1) \xi_j(\mathbf{r}_2) \rangle = g_i \delta_{ij} \delta(\mathbf{r}_2 - \mathbf{r}_1), \quad \langle \xi_i(\mathbf{r}) \rangle = 0, \quad (21)$$

where g_i is proportional to the squared amplitude of the i th interface fluctuation³². This is possible if the distance D between the electron and hole quantum wells is larger than the amplitude of fluctuations on the nearest surfaces.

We consider the characteristic length of the random field potential L to be much shorter than the average distance between excitons $r_s \sim 1/\sqrt{\pi n}$ ($L \ll 1/\sqrt{\pi n}$, where n is the total exciton density). Therefore, in order to obtain the Green's function of the magnetoexcitons with dipole-dipole repulsion in the random field, we obtain the Green's function of a single magnetoexciton in the random field (not interacting with other magnetoexcitons), and then apply the perturbation theory with respect to dipole-dipole repulsion between exciton, using the system of the non-interacting magnetoexcitons as a reference system in analogy with the system without magnetic field³⁰.

Since the effective magnetoexciton Hamiltonian Eq. (17) is translationally invariant, we can write the Green function of isolated magnetoexciton in the momentum space. The Green's function of the center of mass of the isolated magnetoexciton at $T = 0$ in the momentum-frequency domain ($G^{(0)}(\mathbf{p}, \omega)$) in the random field in the coherent potential approximation (CPA) is given by³² (here and below $\hbar = 1$)

$$G^{(0)}(\mathbf{p}, \omega) = \frac{1}{\omega - \varepsilon_0(p) + \mu + iQ(\mathbf{p}, \omega)}, \quad (22)$$

where μ is the chemical potential of the system, and $\varepsilon_0(p) = p^2/2m_H$ is the spectrum of the center mass of the exciton in the “clean” system. The function $Q(\mathbf{p}, \omega)$ is determined by effective random field acting on the center of mass of the exciton. For zero random field,

$Q(\mathbf{p}, \omega) \rightarrow 0$. If $g_i m_H \ll E_b$ the function $Q(\mathbf{p}, \omega)$ in the coherent potential approximation is given by³²

$$Q(\mathbf{p}, \omega) = \frac{1}{2} \int G^{(0)}(\mathbf{q}, \omega) B(|\mathbf{p} - \mathbf{q}|) \frac{d^2 q}{(2\pi)^2}, \quad (23)$$

where

$$B(\mathbf{p}) \equiv \int d^2 R B(\mathbf{R}) e^{-i\mathbf{p}\mathbf{R}}, \quad (24)$$

for $\mathbf{R} = \mathbf{R}_1 - \mathbf{R}_2$, and in the coordinate domain $B(\mathbf{R}_1, \mathbf{R}_2) = \langle V_{eff}(\mathbf{R}_1) V_{eff}(\mathbf{R}_2) \rangle_{av}$ (the effective potential V_{eff} is given by Eq. (18)) has the form³²

$$B(\mathbf{R}_1, \mathbf{R}_2) = B(\mathbf{R}_1 - \mathbf{R}_2) = \frac{\alpha_e^2(g_1 + g_2) + \alpha_h^2(g_3 + g_4)}{2\pi r_H^2} \exp\left(-\frac{(\mathbf{R}_1 - \mathbf{R}_2)^2}{2r_H^2}\right). \quad (25)$$

Note, that, since in the limit of strong magnetic fields the magnetic length $r_H = (eH)^{-1/2}$ is much smaller than the characteristic length of the random field potential L ($r_H \ll L$), and, therefore,

$$\lim_{r_H |\mathbf{R}_1 - \mathbf{R}_2|^{-1} \rightarrow 0} B(\mathbf{R}_1 - \mathbf{R}_2) = \frac{\alpha_e^2(g_1 + g_2) + \alpha_h^2(g_3 + g_4)}{\sqrt{2\pi} r_H} \delta(\mathbf{R}_1 - \mathbf{R}_2), \quad (26)$$

the random field acting on a magnetoexciton is represented by the white noise correlation function of the random potential $B(\mathbf{R}_1, \mathbf{R}_2) = \langle V_{eff}(\mathbf{R}_1) V_{eff}(\mathbf{R}_2) \rangle_{av}$. The time-reversal symmetry of the effective magnetoexciton Schrödinger equation in strong magnetic field, corresponding to the effective Hamiltonian Eq. (8) with the effective random field Eq. (18), and elimination of the long-range property of the random field (Eq. (26)) cancels all effects related with the broken time-reversal symmetry in the Schrödinger equation Eq. (1) at $m_e \neq m_h$, that result in corrections to the Green's function of a magnetoexciton.³⁵

Using Eq. (24), we obtain the Fourier transform of $B(R)$

$$B(p) = \frac{\alpha_e^2(g_1 + g_2) + \alpha_h^2(g_3 + g_4)}{16\pi^4} \exp\left(-\frac{r_H^2 p^2}{32}\right). \quad (27)$$

So the CPA Green's function of the 2D indirect exciton is determined by the solution of the self-consistent equations Eqs. (22) and (23).

In the weak-scattering limit ($g_i m_H \ll E_b \sim e^2/\epsilon D$) we use the second-order Born approximation for Q similar to Refs. [30,31,32], expanding Q (Eq. (23)) in series to the first order in $B(|\mathbf{p} - \mathbf{q}|)$ (which is the first order in g_i), and we replace Eq. (23) by:

$$Q(\mathbf{p}, \omega) = \frac{\pi}{2} \int \delta\left(\omega - \frac{q^2}{2m_H}\right) B(|\mathbf{p} - \mathbf{q}|) \frac{d^2 q}{(2\pi)^2}. \quad (28)$$

Substituting $B(p)$ from Eq. (27) into Eq. (28), we obtain for $Q(\mathbf{p}, \omega)$

$$Q(\mathbf{p}, \omega) = \frac{\alpha_e^2(g_1 + g_2) + \alpha_h^2(g_3 + g_4)}{64\pi^4} m_H \exp\left(-\frac{r_H^2}{32}(p^2 + 2m_H\omega)\right) J_0\left(\frac{r_H^2}{16}\sqrt{2m_H\omega p}\right), \quad (29)$$

where $J_0(z)$ is the Bessel first integral. The second-order Born Green function of the single indirect exciton in the random field in CQW $G^{(0)}(\mathbf{p}, \omega)$ is derived by substituting $Q(\mathbf{p}, \omega)$ from Eq. (29) into Eq. (22).

IV. COLLECTIVE SPECTRUM AND SUPERFLUIDITY OF INDIRECT DIRTY MAGNETOEXCITONS

Due to the interwell separation D , indirect magnetoexcitons both in the ground state and in excited states have electrical dipole moments. We suppose that indirect excitons interact as *parallel* dipoles. This is valid when D is larger than the mean separation $\langle r \rangle$ between electron and hole in the magnetoexciton along quantum wells $D \gg \langle r \rangle$. We take into account that at high magnetic fields $\langle r \rangle \approx Pr_H^2$ ($\langle \mathbf{r} \rangle$ is normal to $\mathbf{P}$) and that the typical value of magnetic momenta (with exactness to logarithm of the exciton density ($\ln(n_{ex})$, see below) is $P \sim \sqrt{n_{ex}}$ (if the dispersion relation $\varepsilon_k(P) = \frac{P^2}{2m_Hk}$ is true). So the inequality $D \gg \langle r \rangle$ is valid at $D \gg \sqrt{n}r_H^2$.

The distinction between excitons and bosons manifests itself in exchange effects (see Refs. [36,16]). These effects for excitons with spatially separated e and h in a rare system $n_{ex}a^2(H, D) \ll 1$ are suppressed due to the negligible overlapping of wave functions of two excitons on account of the potential barrier, associated with the dipole-dipole repulsion of indirect excitons¹⁶ (here n_{ex} , $a(D, H)$ are respectively density and magnetoexciton radius along quantum wells, respectively). The small tunnelling parameter connected with this barrier is $\exp[-\int_{a(H,D)}^{r_0} \sqrt{2m_Hk(\frac{e^2D^2}{R^3} - \frac{\kappa^2}{2m_Hk})}dR]$, where $\kappa^2 \sim \frac{n}{\ln\left(\frac{1}{8\pi nm_H^2 e^4 D^4}\right)}$ is the characteristic momentum of the system (see below); $r_0 = (2m_Hk e^2 D^2 / \kappa^2)^{1/3}$ is the classical turning point for the dipole-dipole potential at the energy equal to the chemical potential of the system. In high magnetic fields the small parameter mentioned above has the form $\exp[-2(m_Hk)^{1/2} e D a^{-1/2}(H, D)]$. Therefore the system of indirect magnetoexcitons can be treated by the diagram technique for a boson system.

The rare excitonic system is stable at $D < D_{cr}(H)$ and $T = 0$ when the magnetoexciton energy $E_b(D, H)$ (calculated in Ref.[24]) is larger than the sum of activation energies $E_L = k \frac{e^2}{\epsilon r_H}$ for incompressible Laughlin liquids of electrons or holes; $k = 0.06$ for $\nu = \frac{1}{3}$ etc.³⁷. Since $k \ll 1$, the critical value $D_{cr} \gg r_H$. In this case one has $E_b = \frac{e^2}{\epsilon D}(1 - \frac{r_H^2}{D^2})$ for a magnetoexciton with quantum numbers $n = m = 0$ (see Ref.[24]). As a result we have from the stability condition (see above) $D_{cr} = r_H(\frac{1}{2k} - 2k)$. For greater ν it gives an upper bound on D_{cr} . In the rare system in high magnetic fields $\mu \ll E_b$ (see below). The coefficient k in the activation energy E_L may be represented as $k = k_0\sqrt{\nu}$. So from the relation between D_{cr} and r_H one has: $\nu_{cr} = \frac{1}{k_0^2} \frac{r_H^2}{D^2}(1 - \frac{r_H^2}{8D^2})$. Thus maximal density for stable magnetoexciton phase is $n_{max} = \nu_{cr}/2\pi r_H^2$.

At $\nu = \frac{1}{2}$ the electron-hole phase can be unstable due to the pairing of electron and hole composite fermions (which form the Fermi surface of composite fermions in the mean field approximation³⁸).

Since the effective Hamiltonian $\hat{H}_{eff}$ (Eq. (17)) of the system of indirect “dirty” magnetoexcitons at small momenta is exactly identical to the Hamiltonian of indirect “dirty” excitons without magnetic field replacing the excitonic mass $M = m_e + m_h$ by the magnetic mass m_H and using the effective random field V_{eff} , we can use the expressions for the ladder approximation Green’s function^{39,40}, collective spectrum, normal and superfluid density and the temperature of Kosterlitz-Thouless phase transition²⁸ for the “dirty” system without magnetic field³⁰ replacing excitonic mass and Q . Since the characteristic length of the random field potential L is much shorter than the average distance between electron and hole, the first step in the averaging procedure with respect to the random field results in the Green function of a non-interacting exciton with the imaginary part. And the second step takes care of the exciton-exciton repulsion. The results of Ref. [30] show the density of the superfluid component and the temperature of the phase transition decrease with the rise of the disorder, which give the good correspondence with the results of the Lopatin-Vinokur approach for the weakly interacting bosons in the presence of disorder.⁴³

Since the characteristic frequencies and momenta which give the greatest contribution to the Green’s function in the ladder approximation are⁴⁰ $\omega \epsilon D / e^2 \sim n / \log[r_H^2 / (nD^4)] \ll 1$ and $pr_H \sim m_H \sqrt{n / \log[r_H^2 / (nD^4)]} \ll 1$, respectively, for the single exciton Green’s function $G^{(0)}(\mathbf{p}, \omega)$, participating in the ladder approximation, we approximate $Q(\mathbf{p}, \omega)$ by $Q(\mathbf{p} =$

$\mathbf{0}, \omega = 0$) (see Eq. (29))

$$Q(\mathbf{p}, \omega) = Q = \frac{\alpha_e^2(g_1 + g_2) + \alpha_h^2(g_3 + g_4)}{64\pi^4} m_H. \quad (30)$$

The chemical potential μ is obtained in the form (compare to Ref.[30]):

$$\mu = \frac{\kappa^2}{2m_H} = \frac{8\pi n}{2m_H \log\left(\frac{\epsilon^2}{8\pi n m_H^2 e^4 D^4}\right)}. \quad (31)$$

where κ is a characteristic momentum. We have the condensate Green's function $D(\mathbf{p}, i\omega_k)$ (compare to Ref.[30]):

$$D^{(0)}(\mathbf{p}, i\omega_k) = \frac{-i(2\pi)^2 n_0 \delta(\mathbf{p})}{i\omega_k + \mu + iQ}, \quad (32)$$

where n_0 is the density of Bose condensate. Since at small temperatures $(n - n_0)/n \ll 1$, according to the ladder approximation³⁹ we use below n instead of n_0 . $G(\mathbf{p}, i\omega_k)$ and $F(\mathbf{p}, i\omega_k)$ are the normal and anomalous Green functions of the overcondensate:

$$\begin{aligned} G(\mathbf{p}, i\omega_k) &= -\frac{i\omega_k + \varepsilon_0(p) + \mu + iQ}{\omega_k^2 + \varepsilon^2(p) - 2i(\mu - \varepsilon_0(p))Q}; \\ F(\mathbf{p}, i\omega_k) &= -\frac{\mu}{\omega_k^2 + \varepsilon^2(p) - 2i(\mu - \varepsilon_0(p))Q}, \end{aligned} \quad (33)$$

where $\varepsilon_0(p)$ is the spectrum of noninteracting excitons; the spectrum of interacting excitons has the form $\varepsilon(p) = \sqrt{\left(p^2/(2m_H) + \sqrt{\mu^2 - Q^2}\right)^2 - (\mu^2 - Q^2)}$, and for small momenta $p \ll \sqrt{2m_H\mu}$ the excitation spectrum is acoustic $\varepsilon(p) = c_s p$, where $c_s = \sqrt{\sqrt{\mu^2 - Q^2}/m_H}$ is the velocity of sound.

At large magnetic momenta P the isolated magnetoexciton spectrum $\varepsilon(P)$ contrary to the case $H = 0$ has a constant limit (being equal to Landau level $\frac{\omega_c}{2}$ for reduced effective mass, see Refs.[24], [18]). As a result the spectrum of the interacting magnetoexciton system also has a plateau at high momenta. Therefore, the Landau criterium of superfluidity is not strictly valid at large P for the interacting magnetoexciton system. However, the probability of excitation of quasiparticles with magnetic momenta $P \gg 1/r_H$ by moving magnetoexciton system is negligibly small at small superfluid velocities²⁵. In this sense, the appearance of a linear branch can be taken as the criterion for superfluidity of 2D magnetoexcitons.

The density of the superfluid component $n_s(T)$ can be obtained using the relation $n_s(T) = n - n_n(T)$, where $n_n(T)$ is the density of the normal component. The density of normal

component n_n is (compare to³⁰):

$$n_n = n_n^0 + \frac{N}{m_H} \int \frac{d\mathbf{p}}{(2\pi)^2} p^2 \mu \frac{\varepsilon_0(p)}{\varepsilon^4(p)} Q. \quad (34)$$

Here N is the total number of particles, and n_n^0 is the density of the normal component in a pure system with no impurities:

$$n_n^0 = -\frac{1}{2m_H} \int \frac{d\mathbf{p}}{(2\pi)^2} p^2 \frac{\partial n_0(p)}{\partial \varepsilon}. \quad (35)$$

where $n_0(p) = (e^{\varepsilon(p)/T} - 1)^{-1}$ is the distribution of an ideal Bose gas of temperature excitations.

The first term in Eq. (34), which does not depend on Q , is the contribution to the normal component due to scattering of quasiparticles with an acoustic spectrum in a pure system at $T \neq 0$ (compare with Ref. [30]),

$$n_n^0 = \frac{3\zeta(3)}{2\pi} \frac{T^3}{c_s^4(n, Q)m_H}, \quad (36)$$

where $\zeta(z)$ is the Riemann zeta function ($\zeta(3) \simeq 1.202$). The second term in Eq. (34) is the contribution to the normal component due to the interaction of the particles (magnetoexcitons) with the random field,

$$n_n = n_n^0 + \frac{nQ}{2m_H c_s^2(n, Q)} = \frac{3\zeta(3)}{2\pi} \frac{T^3}{c_s^4(n, Q)m_H} + \frac{nQ}{2m_H c_s^2(n, Q)}. \quad (37)$$

The density of the superfluid component is $n_s = n - n_n$. From Eqs. (36) and (37) we can see, that the random field decreases the density of the superfluid component.

In a 2D system, superfluidity appears below the Kosterlitz-Thouless transition temperature $T_c = \pi n_s / (2m_H)^{28}$, where only coupled vortices are present. Using the expressions (36) and (37) for the density n_s of the superfluid component, we obtain an equation for the Kosterlitz-Thouless transition temperature T_c . Its solution is

$$T_c = \left[\left(1 + \sqrt{\frac{32}{27} \left(\frac{m_H T_c^0}{\pi n'} \right)^3 + 1} \right)^{1/3} - \left(\sqrt{\frac{32}{27} \left(\frac{m_H T_c^0}{\pi n'} \right)^3 + 1} - 1 \right)^{1/3} \right] \frac{T_c^0}{2^{1/3}}. \quad (38)$$

Here T_c^0 is an auxiliary quantity, equal to the temperature at which the superfluid density vanishes in the mean-field approximation $n_s(T_c^0) = 0$,

$$T_c^0 = \left(\frac{2\pi n' c_s^4 m_H}{3\zeta(3)} \right)^{1/3}. \quad (39)$$

In Eqs. (38) and (39), n' is

$$n' = n - \frac{nQ}{2m_H c_s^2}. \quad (40)$$

Even though the expression similar to Eq. (38) has been already visualized earlier in Ref. [30], the result of this paper is nontrivial because it takes into account the influence of the strong magnetic field on the excitonic superfluidity using mapping of the system with magnetic field onto the system without field by replacing excitonic mass by magnetic mass.

V. DISCUSSION

The superfluid density n_s and the temperature of the Kosterlitz-Thouless transition T_c to the superfluid state for the “pure” system at fixed exciton density n decrease as the magnetic mass m_H increases²⁵ (we can see that by putting $Q = 0$, which makes in Ref. (38) $n' = n$ and $c_s = \sqrt{\mu/m_H}$). Since in strong magnetic fields at $D \gg r_H$ the exciton magnetic mass is $m_H \approx D^3\epsilon/(e^2 r_H^4)$ ²⁴, the superfluid density n_s and the temperature of the Kosterlitz-Thouless transition T_c decrease with increase of the magnetic field H , the parameters of random field α_e , α_h and g_i (Figs. 1 and 2). Since in the “dirty” systems, n_s and T_c decrease with the increase of effective random field Q (analogous to the case without magnetic field Ref.[30]), and in the strong magnetic field Q is proportional to m_H (Eq. (30)), the increase of the magnetic field H increases the effective renormalized random field Q , and suppress the superfluid density n_s and the temperature of the Kosterlitz-Thouless transition T_c .

Note that in the presence of the disorder there is a quantum transition to the superfluid state at zero temperature $T = 0$ with respect to the magnetic field H and the parameters of the disorder α_e , α_h and g_i (Figs. 1 and 2). While in the “pure” system at any magnetic field H at there is always the region in the density-temperature space, where the superfluidity occurs²⁵, in the presence of the disorder at sufficiently large magnetic field H or the parameters of the disorder α_e , α_h and g_i there is no superfluidity at any exciton density.

Note also that in magnetic field the superfluid density n_s and the temperature of the Kosterlitz-Thouless transition T_c decrease when the separation between quantum wells D increases because n_s and T_c are decreasing functions of the magnetoexciton effective mass m_H (Eqs. (37) and (38)), which is an increasing function of D ²⁴. The dependence of n_s and T_c on D through magnetic mass $m_H \approx D^3\epsilon/(e^2 r_H^4)$ and through decreasing dependence on

the random field Q , which increases with the rise of m_H (Eq. (30)), is stronger than their increasing logarithmic dependence on D through the velocity of sound related to the logarithmic dependence on D of the chemical potential of the dipole-dipole repulsion (Eq. (31)). The latter results in the increasing dependence of n_s and T_c on D in a case without magnetic field.³⁰ The decreasing dependence of n_s and T_c on D at high magnetic fields H is significant when D is large ($D \sim 30r_H$), which is far from D used in experiments, because at very large values of D the superfluid system of magnetoexcitons must transform into the system of two incompressible liquids.²³

In the high magnetic field limit at high D , the effective random field is not small, and the approaches assuming coupling with the random field to be much smaller than the dipole-dipole repulsion^{29,41,42,43} are not applicable. Note that in the present work the parameter Q/μ is not required to be sufficiently small, and our formulas for the superfluid density and Kosterlitz-Thouless temperature can be used in the regime of realistic experimental parameters taken from photoluminescence line broadening measurements⁸. The coherent potential approximation (CPA) allowed us to derive the exciton Green's function for the wide range of the random field, and in the weak-scattering limit CPA results in the second-order Born approximation.

The system of “dirty” indirect magnetoexcitons can appear also in unbalanced two-layer *electron* system in CQW in strong magnetic fields near the filling factor $\nu = 1$ ²⁵. An external electric voltage between quantum wells change the filling, so say in the first quantum well the filling factor will be $\nu_1 = \Delta\nu \ll \frac{1}{2}$ and in another one it will be $\nu_2 = 1 - \Delta\nu$. Unbalanced filling factors $\nu_1 = 1 + \Delta\nu$, $\nu_2 = 1 - \Delta\nu$ is also possible. Thus in the first quantum well (QW) there are rare electrons on the second Landau level, and in the second QW there are rare empty places (holes) in the first Landau level. Excess electrons in the first QW and holes in the second QW can bind to indirect magnetoexcitons with the density $n_{ex} = eH\Delta\nu/2\pi$. Superfluidity in two-layer $e - e$ system in high magnetic fields in the cases mentioned is analogous to the superfluidity of two-layer $e - h$ system²⁵.

Note that in this paper we assumed spin degeneracy factor $s = 1$ in high magnetic fields H . Since at $H = 0$ the spin degeneracy is $s = 4$,³⁰ at low H lifting spin degeneracy will have substantial effect on the magnetoexciton superfluidity, which we do not take into account in this paper where we consider only high H .

Acknowledgements

We thank Drs. Valerii Vinokur, Alexei Lopatin (Argonne National Laboratory), Profs. Peter Littlewood and Roland Zimmermann and the participants of 1st Int. Conf. on Spontaneous Coherence in Excitonic Systems for many useful and stimulating discussions. Yu. E. L. was supported by the INTAS and RFBR grants. D. W. S. and R. D. C. have been supported by the National Science Foundation.

-
- ¹ L. V. Butov, A. Zrenner, G. Abstreiter, G. Bohm and G. Weimann, Phys. Rev. Lett. **73**, 304 (1994); L. V. Butov and A. I. Filin, Phys. Rev. **B 58**, 1980 (1998); L. V. Butov, C. W. Lai, A. L. Ivanov, A. C. Gossard, and D. S. Chemla, Nature **417**, 47 (2002); L. V. Butov, A. C. Gossard, and D. S. Chemla, Nature **418**, 751 (2002)).
- ² S. Denev, V. Negoita, D. W. Snoke, B. Laikhtman, K. Eberl, and L. Pfeiffer, Phys. Rev. **B 66**, 205304 (2002).
- ³ V. V. Krivolapchuk, E. S. Moskalenko, and A. L. Zhmodikov, Phys. Rev. **B 64**, 045313 (2001).
- ⁴ A. V. Larionov, V. B. Timofeev, J. Hvam, and K. Soerensen, ZhETF **117**, 1255 (2000) [JETP **90**, 1093 (2000)]; A. V. Larionov and V. B. Timofeev, Pis'ma ZhETF **73**, 342 (2001) [JETP Lett. **73**, 301 (2001)]; A. V. Larionov, V. B. Timofeev, J. Hvam, and K. Soerensen, Pis'ma ZhETF **75**, 233 (2002) [JETP Lett. **75**, 200 (2002)].
- ⁵ T. Fukuzawa, E. E. Mendez and J. M. Hong, Phys. Rev. Lett. **64**, 3066 (1990); J. A. Kash, M. Zachau, E. E. Mendez, J. M. Hong and T. Fukuzawa, Phys. Rev. Lett., **66**, 2247 (1991).
- ⁶ U. Sivan, P. M. Solomon and H. Strikman, Phys. Rev. Lett. **68**, 1196 (1992).
- ⁷ For a general review of experiments in quantum wells, see chapter 10 of S. A. Moskalenko and D. W. Snoke, *Bose-Einstein Condensation of Excitons and Biexcitons and Coherent Nonlinear Optics with Excitons* (Cambridge University Press, New York, 2000).
- ⁸ D. Snoke, S. Denev, Y. Liu, L. Pfeiffer and K. West, Nature **418**, 754 (2002).
- ⁹ D. Snoke, Science **298**, 1368 (2002).
- ¹⁰ Yu. E. Lozovik and V. I. Yudson, Pis'ma ZhETF **22**, 26 (1975) [JETP Lett. **22**, 26(1975)]; ZhETF **71**, 738 (1976) [JETP **44**, 389 (1976)]; Sol. St. Comms. **18**, 628 (1976); Sol. St. Comms. **18**, 628 (1976); Sol. St. Comms. **21**, 211 (1977).

- ¹¹ A. V. Klyuchnik and Yu. E. Lozovik, ZhETF **76**, 670 (1979) [JETP **49**, 335 (1979)]; Yu. E. Lozovik and A. V. Poushnov, Phys. Lett. **A 228**, 399 (1997); .
- ¹² S. I. Shevchenko, Phys. Rev. Lett. **72**, 3242 (1994).
- ¹³ J. Zang, D. Schmeltzer and J. L. Birman, Phys. Rev. Lett. **71**, 773 (1993).
- ¹⁴ Xu. Zhu, P. B. Littlewood, M. S. Hybertsen and T. M. Rice, Phys. Rev. Lett. **74**, 1633 (1995).
- ¹⁵ S. Conti, G. Vignale and A. H. MacDonald, Phys. Rev. **B 57**, R6846 (1998).
- ¹⁶ Yu. E. Lozovik and O. L. Berman, Pis'ma ZhETF **64**, 526 (1996) [JETP Lett. **64**, 573 (1996)]; ZhETF **111**, 1879 (1997) [JETP **84**, 1027 (1997)].
- ¹⁷ Yu. E. Lozovik, O. L. Berman, and M. Willander, J. Phys.: Condens. Matter **14**, 12457 (2002).
- ¹⁸ I. V. Lerner and Yu. E. Lozovik, ZhETF **74**, 274 (1978) [JETP **47**, 146 (1978)]; ZhETF **76**, 1136 (1979) [JETP **49**, 576 (1979)]; ZhETF **78**, 1167 (1980) [JETP **51**, 588 (1980)]; ZhETF **80**, 1488 (1981) [JETP **53**, 763 (1981)];
- ¹⁹ A. B. Dzjubenko and Yu. E. Lozovik, Fiz. Tverd. Tela **25**, 1519 (1983) [Solid State Phys. **25**, 874 (1983)]; Fiz. Tverd. Tela **26**, 1540 (1984) [Solid State Phys. **26**, 938 (1984)]; J. Phys. **A 24**, 415 (1991); Yu. E. Lozovik, I. V. Ovchinnikov, S. Yu. Volkov, L. V. Butov, and D. S. Chemla, Phys. Rev. **B 65**, 030223 (2002).
- ²⁰ D. Paquet, T. M. Rice, and K. Ueda, Phys. Rev. **B 32**, 5208 (1985).
- ²¹ C. Kallin and B. I. Halperin, Phys. Rev. **B 30**, 5655 (1984); Phys. Rev. **B 31**, 3635 (1985).
- ²² D. S. Chemla, J. B. Stark, and W. H. Knox, In *"Ultrafast Phenomena VIII"*, Eds. J. -L. Martin et al., Springer, 21 (1993); C. W. Lai, J. Zoch, A. C. Gossard, and D. S. Chemla, Science **303**, 503 (2004).
- ²³ D. Yoshioka and A. H. MacDonald, J. Phys. Soc. of Japan, **59**, 4211 (1990).
- ²⁴ Yu. E. Lozovik and A. M. Ruvinsky, Phys. Lett. **A 227**, 271 (1997); ZhETF **112**, 1791 (1997) [JETP **85**, 979 (1997)].
- ²⁵ Yu. E. Lozovik, O. L. Berman and V. G. Tsvetus. Pis'ma ZhETF **66**, 332 (1997); Phys. Rev. **B 59**, 5627 (1999).
- ²⁶ M. A. Olivares-Robles and S. E. Ulloa, Phys. Rev. **B 64**, 115302 (2001).
- ²⁷ S. A. Moskalenko, M. A. Liberman, D. W. Snoke and V. V. Botan, Phys. Rev. **B 66**, 245316 (2002).
- ²⁸ J. M. Kosterlitz and D. J. Thouless, J. Phys. **C 6**, 1181 (1973); D. R. Nelson and J. M. Kosterlitz, Phys. Rev. Lett. **39**, 1201 (1977).

- ²⁹ Yu. E. Lozovik, O. L. Berman and A. M. Ruvinsky, Pisma ZhETF **69**, 573 (1999) [JETP Lett. **69**, 616 (1999)].
- ³⁰ O. L. Berman, Yu. E. Lozovik, D. W. Snoke and R. D. Coalson, Phys. Rev. B **70**, 235310 (2004).
- ³¹ Zh. S. Gevorkyan and Yu. E. Lozovik, Fiz. Tverd. Tela **27**, 1800 (1985) [Sov. Phys. Solid State **27**, 1079 (1985)].
- ³² Yu. E. Lozovik and A. M. Ruvinsky, ZhETF **114**, 1451 (1998) [JETP **87**, 788 (1998)].
- ³³ L. P. Gor'kov and I. E. Dzyaloshinskii, ZhETF, **53**, 717 (1967) [JETP **26**, 449 (1968)].
- ³⁴ V. L. Bonch-Bruевич, I. P. Zvyagin, R. Kaïper, A. G. Mironov, R. Énderlaïn, and B. Ésser, in *Electronic Theory of Disordered Semiconductors*, p. 350 , Nauka, Moscow (1981).
- ³⁵ P. I. Arseyev and A. B. Dzyubenko, Phys. Rev. **B 52**, R2261 (1995).
- ³⁶ L. V. Keldysh and A. N. Kozlov, ZhETF, **54**, 978 (1968) [JETP, **27**, 521 (1968)].
- ³⁷ R. B. Laughlin, in *The Quantum Hall Effect*, edited by R. E. Prange and S. M. Girvin (eds.), Springer-Verlag, N.Y. (1987).
- ³⁸ B. I. Halperin, P. A. Lee and N. Read, Phys. Rev. **B 47**, 7312 (1993).
- ³⁹ A. A. Abrikosov, L. P. Gor'kov, and I. E. Dzyaloshinskii, *Methods of Quantum Field Theory in Statistical Physics*, Prentice-Hall, Englewood Cliffs, N.J. (1963).
- ⁴⁰ Yu. E. Lozovik and V. I. Yudson, Physica **A 93**, 493 (1978).
- ⁴¹ K. Huang, in *Bose-Einstein Condensation*, A. Griffin, D. W. Snoke, S. Stringari, Eds. (Cambridge Univ. Press, Cambridge, 1995), p.p. 31-50.
- ⁴² K. Huang and H. F. Meng, Phys. Rev. Lett. **69**, 644 (1992); H. F. Meng, Phys. Rev. **B 49**, 1205 (1994).
- ⁴³ A. V. Lopatin and V. M. Vinokur, Phys. Rev. Lett. **88**, 235503 (2002).




FIG. 1: Dependence of temperature of Kosterlitz-Thouless transition $T_c = T_c(H)$ (in units K ; for $GaAs/GaAsAl$: $\epsilon = 13$ on the magnetic field H (in units T) at $g_i = 10^4 nm^4$ at the interwell distance $D = 15 nm$; at the exciton density $n = 1.0 \times 10^{11} cm^{-2}$; at the different parameters of the random field $\alpha = \alpha_e = \alpha_h$: $\alpha = 0$ – solid curve; $\alpha = 0.5 meV/nm$ – dotted curve; $\alpha = 0.7 meV/nm$ – dashed curve.




FIG. 2: Dependence of temperature of Kosterlitz-Thouless transition $T_c = T_c(n)$ (in units K ; for $GaAs/GaAsAl$: $M = 0.24m_0$; $\epsilon = 13$; m_0 is a mass of electron) on the exciton density n (in units cm^{-2}) at the interwell distance $D = 15nm$ at the parameters of the random field $\alpha_e = \alpha_h = 1.5meV/nm$; $g_i = 10^4 nm^4$; at different magnetic fields H (in units of T): $H = 0$ – solid curve (for $H = 0$ we use the results of Ref.[30] for spin degeneracy factor $s = 4$); $H = 14T$ – dotted curve; $H = 15T$ – dashed curve.